

COMPONENT[®]
BRAND IMPLANTS

VetLife

TE-200 WITH Tylan[®]

FOR STEERS AND HEIFERS

fed in confinement for slaughter

trenbolone acetate 200 mg and estradiol

USP 20 mg with 29 mg tylosin tartrate

as a local antibacterial



20 dose CARTRIDGE BELT*

For Use In Animals Only

Each CARTRIDGE BELT holds 20 doses of COMPONENT[®] TE-200 with Tylan Implants. Each dose of 11 pellets consists of 10 pellets containing a total of 200 mg of trenbolone acetate and 20 mg estradiol plus 1 pellet containing 29 mg tylosin tartrate as a local antibacterial. COMPONENT[®] TE-200 Implants are recommended for use in steers and heifers fed in confinement for slaughter for INCREASED RATE OF WEIGHT GAIN AND IMPROVED FEED EFFICIENCY.

INDICATIONS FOR USE: This product contains trenbolone acetate and estradiol for increased rate of weight gain and improved feed efficiency in a slow-release delivery system. Do not use in veal calves. Effectiveness and animal safety in veal calves have not been established.

NOTE: COMPONENT[®] TE-200 is not better for increased average daily gain or improved feed efficiency than COMPONENT[®] TE-H (140 mg trenbolone acetate / 14 mg estradiol) in heifers fed in confinement for slaughter. Studies have demonstrated that administration of COMPONENT[®] TE-200 can result in decreased marbling scores when compared to non-implanted steers and heifers.

DOSAGE: Administer 1 implant per animal according to the following instructions.

GENERAL INSTRUCTIONS: Study the instructions which should be followed carefully at all times, avoiding short cuts. Skin infection can be avoided by properly preparing implant site and implanter. During fly season use fly repellent on implant site. One designated team member should always do the implanting. Cleanliness of hands and instruments is important at all times.

WARNING

Not to be used in animals intended for subsequent breeding, or in dairy animals. Implant one dose — 11 pellets (entire contents of one cartridge cell) — in the ear subcutaneously. Any other site of implantation is in violation of Federal Law. **DO NOT** attempt to salvage implantation site for animal feed or human food.



A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

**RESTRICTED DRUG — USE ONLY AS DIRECTED (CALIFORNIA) —
FOR USE IN ANIMALS ONLY — NOT FOR HUMAN USE —
KEEP OUT OF REACH OF CHILDREN**

IMPLANTING INSTRUCTIONS:


Loading the Implanter

Load the implanter following the instructions supplied with each implanter.



Restrain the Animal

Speed of implantation as well as safety of handlers is best achieved by restraining animal in a squeeze chute using head restraint. When implanting horned cattle, better control is obtained with additional use of nose tongs.




Prepare the Implant Site

Scrub the back side of the ear (implant site) with a piece of clean absorbent cotton which has been soaked with topical germicidal solution. Follow manufacturer's directions on germicide for correct strength and preparation of solution. Avoid getting disinfectant in animal's eyes.



Where to Implant

The full contents of one cartridge cell should be implanted beneath the skin on the back side of the middle one-third of the ear illustrated in the drawing. The implant must not be closer to the head than the edge of the auricular cartilage ring farthest from the head. The location for insertion of the needle is a point toward the tip of the ear at least a needle length away from the intended deposition site. Avoid injuring the large arteries, veins and cartilage of the ear.



Insert the Needle

With one hand firmly grasp the ear. With the other hand insert needle point through the skin and ease forward on a lateral plane until the entire length of the needle is under the skin.

Implant the Pellets

After inserting the needle fully in the correct implant position, squeeze the trigger fully as the needle is withdrawn from the ear. This properly deposits the implant in the needle track. This procedure should prevent breakage or crushing of pellets if otherwise forced into contact with tough fibrous-tissue underlying the skin. The length and total contact area of a single dose are designed to permit absorption of the hormones after implantation to stimulate good weight gain. Broken or crushed pellets may interfere with rates of gain.

Clean the Needle

Clean the implanter needle with alcohol or a properly diluted germicidal solution prior to implanting the next animal.

Storage Conditions

Store in a refrigerator (2°-8°C; 36°-46°F) and protect from sunlight. Use before the expiration date printed on the foil pouch.

ANADA 200-346, Approved by FDA

Manufactured by a non-sterilizing process.

*U.S. Patent No. 4,531,938

COMPONENT TE-200 with Tylan is covered by U.S. Patent No. 5,874,098

COMPONENT is a registered trademark of Ivy Animal Health, Inc.

Tylan is a registered trademark of Eli Lilly and Company.

Manufactured for **VetLife** by Ivy Laboratories • Overland Park, KS 66214, USA

VetLife and Ivy Laboratories are divisions of Ivy Animal Health, Inc.