

BLUE BIRD TURKEY FEED

TYPE C: MEDICATED FEED

INDICATIONS

For the prevention of coccidiosis caused by *E. adenoeides*, *E. meleagrimitis* and *E. gallopavonis*, and for increased rate of weight gain and improved feed efficiency in growing turkeys.

ACTIVE INGREDIENT

Monensin¹ (as monensin sodium) 54 - 90 g/ton*
Bacitracin Methylene Disalicylate..... 4 - 50 g/ton*

GUARANTEED ANALYSIS

Nutritional Ingredients _____ %

INGREDIENTS

As listed in AAFCO Handbook.

Light mineral oil may be added to control dusting.

DIRECTIONS FOR USE

Feeding Directions: Feed continuously as the sole ration. The optimum level depends upon the severity of coccidiosis exposure.

CAUTION: Do not allow horses, other equines, mature turkeys or guinea fowl access to feed containing monensin. Ingestion of monensin by horses and guinea fowl has been fatal. Some strains of turkey coccidia may be monensin tolerant or resistant. Monensin may interfere with development of immunity to turkey coccidiosis.

Store in a cool, dry place.

MANUFACTURED BY

BLUE BIRD FEED MILL

Anytown, U.S.A. 12345

Net Weight lb (kg) on bag or bulk

Lot No. _____

*The medicated feed label must state a single drug concentration.

¹Sourced from Coban, NADA#038-878

