

Monensin and Bacitracin Methylene Disalicylate Medicated Broiler Chicken Feed Type C Medicated Feed

INDICATIONS

To aid in the prevention of coccidiosis caused by *Eimeria necatrix*, *E. tenella*, *E. acervulina*, *E. brunetti*, *E. mivati*, and *E. maxima*, and for increase in rate of weight gain and improved feed efficiency.

ACTIVE DRUG INGREDIENT

Monensin ³ , USP	90 to 110 g/ton*
Bacitracin Methylene Disalicylate.....	5 to 25 g/ton*

GUARANTEED ANALYSIS

Crude Protein, not less than.....	_____	%
Lysine, not less than.....	_____	%
Methionine, not less than	_____	%
Crude Fat, not less than	_____	%
Crude Fiber, not more than	_____	%
Calcium, not less than.....	_____	%
Calcium, not more than.....	_____	%
Phosphorus, not less than.....	_____	%
Salt ¹ , not less than.....	_____	%
Salt ¹ , not more than.....	_____	%
Sodium ² , not less than	_____	%
Sodium ² , not more than	_____	%

¹If added

²Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

INGREDIENTS

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials.

FEEDING DIRECTIONS

Feed continuously as the sole ration.

CAUTION

Do not feed to laying chickens. The use of monensin with no withdrawal period may limit feed intake resulting in reduced weight gain. Do not allow horses, other equines, mature turkeys, or guinea fowl access to feed containing monensin. Ingestion of monensin by horses and guinea fowl has been fatal.

MANUFACTURED BY

BLUE BIRD FEED MILL

Any town, USA 12345

Net Weight lb (kg) on bag or bulk

*The medicated feed label must state a single drug concentration.

³Sourced from Coban, NADA# 038-878

