

TURKEY RATION TYPE C MEDICATED FEED

Caution: Federal law restricts medicated feed containing this drug to use by or on the order of a licensed veterinarian.

Do Not Feed Undiluted

INDICATIONS

For the prevention of coccidiosis caused by *Eimeria adenoeides*, *E. meleagrimitis*, and *E. gallopavonis*, and for increased rate of weight gain and improved feed efficiency in growing turkeys.

ACTIVE DRUG INGREDIENT

Monensin ³	54 to 90 g/ton*
Virginiamycin	10 to 20 g/ton*

GUARANTEED ANALYSIS

Crude Protein, not less than.....	_____	%
Lysine, not less than.....	_____	%
Methionine, not less than	_____	%
Crude Fat, not less than	_____	%
Crude Fiber, not more than	_____	%
Calcium, not less than.....	_____	%
Calcium, not more than	_____	%
Phosphorus, not less than.....	_____	%
Salt ¹ , not less than.....	_____	%
Salt ¹ , not more than.....	_____	%
Sodium ² , not less than	_____	%
Sodium ² , not more than	_____	%

¹If added.

²Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

INGREDIENTS

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials' Publication (AAFCO Official Publication).

DIRECTIONS FOR USE

Feed continuously as the sole ration from 1 day of age to 10 weeks of age.

CAUTION: Do not allow horses, other equines, mature turkeys or guinea fowl access to feed containing monensin. Ingestion of monensin by horses, mature turkeys, and guinea fowl has been fatal. Some strains of turkey coccidian may be monensin tolerant or resistant. Monensin may interfere with the development of immunity to turkey coccidiosis.

MANUFACTURED BY

Blue Bird Feed Mill
Anytown, USA 12345

Net Weight lb (kg) on bag or bulk

*The medicated feed label must state a single drug concentration.

³Sourced from Coban, NADA#038-878



Elanco, Coban and the diagonal bar are trademarks owned or licensed by Eli Lilly and Company, its subsidiaries or affiliates.

YL090145AMX