

# Blue Bird Broiler Ration NaNi

## Type C Medicated Feed

Do Not Feed Undiluted.

### INDICATIONS

For the prevention of coccidiosis caused by *Eimeria tenella*, *E. necatrix*, *E. acervulina*, *E. maxima*, *E. brunetti*, and *E. mivati*.

### ACTIVE DRUG INGREDIENT

Narasin <sup>1</sup> .....	27 – 45 g/ton*
Nicarbazin <sup>1</sup> .....	27 – 45 g/ton*

### GUARANTEED ANALYSIS

To provide nutritional requirements.

### INGREDIENTS

As listed in AAFCO Handbook.

### MIXING DIRECTIONS

Feed continuously as the only ration.

**WARNING:** Withdraw 5 days before slaughter.

**CAUTION:** Nicarbazin medicated broilers may show reduced heat tolerance if exposed to high temperature and high humidity. Provide adequate drinking water and ventilation during these periods. Do not allow adult turkeys, horses or other equines access to formulations containing narasin. Ingestion of narasin by these species has been fatal. Do not feed to laying hens.

### MANUFACTURED BY

BLUE BIRD FEED MILL

Robin, IN 46813

\*The medicated feed label must state a single drug concentration. Ingredients must be provided in 1:1 ratio.

<sup>1</sup>Sourced from Maxiban, NADA#138-952



Elanco, Maxiban and the diagonal bar are trademarks owned or licensed by Eli Lilly and Company, its subsidiaries or affiliates.

YL090146AMB