

# Monensin Medicated Goat Feed

## Type B Medicated Feed

### FEED ONLY TO GOATS MAINTAINED IN CONFINEMENT

For the prevention of coccidiosis caused by *Eimeria crandallis*, *E. christensenii*, and *E. ninakohlyakimovae* in goats maintained in confinement

#### Active Drug Ingredient

Monensin, USP<sup>6</sup> .....21-80,000 g/ton\*

#### Guaranteed Analysis

Crude Protein, not less than.....	_____	%
Non-Protein Nitrogen (NPN) <sup>1</sup> , not more than .....	_____	%
Crude Fat, not less than .....	_____	%
Crude Fiber, not more than .....	_____	%
Calcium, not less than.....	_____	%
Calcium, not more than .....	_____	%
Phosphorus, not less than.....	_____	%
Salt <sup>2</sup> , not less than.....	_____	%
Salt <sup>2</sup> , not more than.....	_____	%
Sodium <sup>3</sup> , not less than .....	_____	%
Sodium <sup>3</sup> , not more than .....	_____	%
Copper <sup>2,4</sup> , not less than .....	_____	ppm
Copper <sup>2,4</sup> , not more than .....	_____	ppm
Selenium, not less than .....	_____	ppm
Vitamin A <sup>2,5</sup> , not less than.....	_____	I.U./lb

<sup>1</sup>When added.

<sup>2</sup>If added.

<sup>3</sup>Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

<sup>4</sup>If total copper exceeds 20 ppm.

<sup>5</sup>Other than precursors of Vitamin A.

#### Ingredients

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials.

---

\*Final printed label on formulated Type B medicated feed must bear a single drug concentration.

## Mixing Directions

Thoroughly mix monensin Type B Medicated Feed into the appropriate amount of non-medicated feed to obtain the correct concentration in the Type C Medicated Feed, 20 g/ton, as follows:

<b>Monensin Level Desired in Type C Medicated Feed</b>	<b>Monensin Level in Type B Medicated Feed</b>	<b>Amount of Type B Medicated Feed to make a ton of Type C Medicated Feed</b>	<b>Amount of Non-Medicated Feed</b>
<b>g/ton</b>	<b>g/ton</b>	<b>lb/ton</b>	<b>lb/ton</b>
20	100	400	1600
20	300	133	1867
20	600	67	1933
20	1200	33	1967

## Caution

Do not allow horses or other equines access to feed containing monensin. Ingestion of monensin by horses has been fatal. Monensin medicated cattle and goat feeds are safe for use in cattle and goats only. Consumption by unapproved species may result in toxic reactions. Feeding undiluted or mixing errors resulting in high concentrations of monensin has been fatal to cattle and could be fatal to goats. Must be thoroughly mixed in feeds before use. Do not feed undiluted. Do not exceed the levels of monensin recommended in the feeding directions, as reduced average daily gains may result. Do not feed to lactating goats.

## Manufactured By

Blue Bird Feed Mill  
Anytown, USA 12345

Net Weight lb on Bag or Bulk

<sup>6</sup>Sourced from Rumensin, NADA#095-735

