

MONENSIN TYPE B MEDICATED FEED FOR MATURE REPRODUCING BEEF COWS (ON PASTURE OR IN DRY LOT)

For improved feed efficiency when receiving supplemental feed;
For prevention and control of coccidiosis caused by *Eimeria bovis* and *Eimeria zuernii*.

Active Drug Ingredient

Monensin, USP⁵401-80,000 g/ton*

Guaranteed Analysis

Crude Protein, not less than.....	_____ %
Non-Protein Nitrogen (NPN) ¹ , not more than	_____ %
Crude Fat, not less than	_____ %
Crude Fiber, not more than	_____ %
Calcium, not less than	_____ %
Calcium, not more than	_____ %
Phosphorus, not less than.....	_____ %
Salt ² , not less than.....	_____ %
Salt ² , not more than.....	_____ %
Sodium ³ , not less than	_____ %
Sodium ³ , not more than	_____ %
Potassium, not less than	_____ %
Vitamin A ^{2,4} , not less than.....	_____ I.U./lb

¹When added.

²If added.

³Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

⁴Other than precursors of Vitamin A.

Ingredients

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials.

*Final printed label on formulated Type B medicated feed must bear a single drug concentration

Mixing Directions

Thoroughly mix monensin Type B Medicated Feed into the appropriate amount of non-medicated feed to obtain the correct concentration in the Type C Medicated Feed, 25 to 400 g/ton. Use only the portion of the table below that is applicable to the concentration of monensin in the Type C Medicated Feed you manufacture.

Monensin Level Desired in Type C Medicated Feed	Monensin Level in Type B Medicated Feed	Amount of Type B Medicated Feed to make a ton of Type C Medicated Feed	Amount of Non-Medicated Feed
g/ton (mg/lb)	g/ton	lb/ton	lb/ton
50 (25)	500	200.0	1800
100 (50)	600	333.4	1666
200 (100)	800	500.0	1500
300 (150)	1000	600.0	1400
400 (200)	1200	666.6	1333

Caution

Do not allow horses or other equines access to feed containing monensin. Ingestion of monensin by horses has been fatal. Monensin medicated cattle and goat feeds are safe for use in cattle and goats only. Consumption by unapproved species may result in toxic reactions. Feeding undiluted or mixing errors resulting in high concentrations of monensin has been fatal to cattle and could be fatal to goats. Must be thoroughly mixed in feeds before use. Do not feed undiluted. Do not exceed the levels of monensin recommended in the feeding directions, as reduced average daily gains may result. If feed refusals containing monensin are fed to other groups of cattle, the concentration of monensin in the refusals and amount of refusals fed should be taken into consideration to prevent monensin overdosing.

Warning

A withdrawal period has not been established for this product in preruminating calves. Do not use in calves to be processed for veal.

Manufactured By

Blue Bird Feed Mill
Anytown, USA 12345

Net Weight lb on Bag or Bulk

⁵Sourced from Rumensin, NADA#095-735

