

# Tilmicosin Medicated Cattle Feed

## Type C Medicated Feed

**CAUTION:** Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

### INDICATION

For the control of bovine respiratory disease (BRD) associated with *Mannheimia haemolytica*, *Pasteurella multocida*, and *Histophilus somni* in groups of beef and non-lactating dairy cattle, where active BRD has been diagnosed in at least 10% of the animals in the group.

### ACTIVE DRUG INGREDIENT

Tilmicosin (as tilmicosin phosphate)<sup>a</sup> ..... 568 to 757 grams per ton<sup>\*, b</sup>.

### GUARANTEED ANALYSIS

Crude Protein, not less than .....	_____	%
Non-Protein Nitrogen (NPN) <sup>1</sup> , not more than .....	_____	%
Crude Fat, not less than .....	_____	%
Crude Fiber, not more than .....	_____	%
Calcium, not less than .....	_____	%
Calcium, not more than .....	_____	%
Phosphorus, not less than .....	_____	%
Salt <sup>2</sup> , not less than .....	_____	%
Salt <sup>2</sup> , not more than .....	_____	%
Sodium <sup>3</sup> , not less than .....	_____	%
Sodium <sup>3</sup> , not more than .....	_____	%
Potassium, not less than .....	_____	%
Vitamin A <sup>2, 4</sup> , not less than .....	_____	I.U./lb

<sup>1</sup> When added.

<sup>2</sup> If added.

<sup>3</sup> Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

<sup>4</sup> Other than precursors of Vitamin A.

### INGREDIENTS

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials (AAFCO).

---

### FEEDING DIRECTIONS

This tilmicosin feed is to be fed continuously for a single, 14-day period at 568 grams to 757 grams (626 ppm to 834 ppm) per ton on a 100% dry matter basis of Type C medicated feed as the sole ration to provide 12.5 mg tilmicosin/kg of body weight/day.

Complete Type C medicated feeds containing tilmicosin should not be pelleted. Do not use in any feeds containing bentonite, cottonseed meal, or cottonseed hulls. Bentonite, cottonseed meal, or cottonseed hulls in feeds may affect the efficacy of tilmicosin.

### CAUTION:

Use only in cattle fed in confinement for slaughter.

Do not allow horses or other equines access to feeds containing tilmicosin. The safety of tilmicosin has not been established in cattle intended for breeding purposes.

To assure both food safety and responsible use in cattle, the treatment of cattle with this medicated feed is required to be initiated within the first 45 days of the production period. The treatment should not occur concurrent with or following administration of an injectable macrolide, or within 3 days following administration of a non-macrolide injectable BRD therapy

Tilmicosin medicated feed treatment has not been evaluated in cattle with severe clinical disease. Cattle with severe clinical illness should be evaluated for individual treatment with an alternative non-macrolide therapy.

The expiration date for a tilmicosin Veterinary Feed Directive (VFD) for cattle must not exceed 45 days from the time of issuance. VFDs for tilmicosin phosphate shall not be refilled.

**WARNINGS:**

**RESIDUE WARNING:** Cattle intended for human consumption must not be slaughtered within 28 days of the last treatment with this drug product.

This drug product is not approved for use in female dairy cattle 20 months of age or older. Use in these cattle may cause drug residues in milk. This drug product is not approved for use in calves intended to be processed for veal. A withdrawal period has not been established in pre-ruminating calves.

**MANUFACTURED BY**  
**Bluebird Feed Company**  
**Any town, USA 12345**

**Net Weight:** \_\_\_\_\_ lb ( \_\_\_\_\_ kg)

For emergency medical information, to report an adverse effect, or for technical service call: 1-800-428-4441.

<sup>a</sup> Sourced from Pulmotil, NADA# 141-064

\* Final printed label must bear a single drug concentration.

<sup>b</sup> 100% dry matter basis

Elanco, Pulmotil and the diagonal bar are trademarks owned or licensed by Eli Lilly and Company, its subsidiaries or affiliates.

