

Net Weight on bag or bulk

FREE CHOICE MONENSIN MINERAL GRANULES TYPE C MEDICATED FEED

For increased rate of weight gain; For prevention and control of coccidiosis caused by *Eimeria bovis* and *Eimeria zuernii* in growing cattle on pasture or in dry lot (stocker and feeder cattle and dairy and beef replacement heifers).

ACTIVE DRUG INGREDIENT

Monensin sodium¹ 1,620 g/ton

GUARANTEED ANALYSIS

Salt, not more than.....	26.0 %
Salt, not less than.....	22.0 %
Calcium, not more than.....	10.5 %
Calcium, not less than.....	9.0 %
Phosphorus, not less than.....	6.0 %
Magnesium, not less than	_____ %
Potassium, not less than	_____ %
Sodium ^a , not more than.....	_____ %
Sodium ^a , not less than	_____ %
Copper, not less than	_____ ppm
Selenium, not less than	_____ ppm
Zinc, not less than	_____ ppm
Vitamin A, not less than	_____ IU/lb

^a Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

INGREDIENTS

Salt, dried cane molasses, monocalcium phosphate, ground limestone or calcium carbonate, dicalcium phosphate, processed grain by-products, cane molasses, and mineral oil (include microingredients as appropriate). Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials.

FEEDING DIRECTIONS

For free choice feeding to growing cattle on pasture or in dry lot (stocker and feeder cattle and dairy and beef replacement heifers). Feed continuously on a free-choice basis at the rate of 50 to 200 milligrams per head per day. Place in covered mineral feeders located near the animals' water supply and/or loafing area. Provide one feeder for each 20 head. Fill feeders with a quantity of Monensin Mineral Supplement that will be consumed in

seven days. It is essential to offer enough feeding stations to insure that all animals have free access at all times to the Monensin Mineral Supplement. Cattle should consume 1 to 4 ounces per head daily which will provide the approved effective intake of 50 to 200 mg monensin.

Note: Adequate consumption for cattle on pasture is dependent on good pasture conditions.

CAUTIONS

During the first 5 days of feeding, cattle should receive no more than 100 mg/hd/day. Do not feed additional salt or minerals. Do not mix with grain or other feeds. Stressed and/or water deprived cattle should be adapted to the pasture and to unmedicated mineral supplement before using the Monensin Mineral Supplement. Do not allow horses or other equines access to feeds containing monensin. Ingestion of monensin by horses has been fatal. Monensin medicated cattle feeds are safe for use in cattle only. Consumption by unapproved species may result in toxic reactions. Do not exceed the levels of monensin recommended in the feeding directions as reduced average daily gains may result. Do not feed to lactating goats.

WARNING

A withdrawal period has not been established for monensin in pre-ruminating calves. Do not use in calves to be processed for veal.

Manufactured By

Blue Bird Feed Mill

Anytown, USA 12345

¹Sourced from Rumensin, NADA#095-735

