

Tylosin Medicated Feed – Swine Ileitis with Tylosin in Drinking Water

Type C Medicated Feed

CAUTION: Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

INDICATION

For control of porcine proliferative enteropathies (PPE, ileitis) associated with *Lawsonia intracellularis* immediately after medicating with tylosin in drinking water.

ACTIVE DRUG INGREDIENT

Tylosin (as tylosin phosphate)^a 40 to 100 g/ton*

GUARANTEED ANALYSIS

Crude Protein, not less than.....	_____	%
Lysine, not less than.....	_____	%
Crude Fat, not less than	_____	%
Crude Fiber, not more than	_____	%
Calcium, not less than.....	_____	%
Calcium, not more than	_____	%
Phosphorus, not less than.....	_____	%
Salt ¹ , not less than.....	_____	%
Salt ¹ , not more than.....	_____	%
Sodium ² , not less than	_____	%
Sodium ² , not more than	_____	%
Selenium, not less than	_____	ppm
Zinc, not less than	_____	ppm

¹ If added.

² Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

INGREDIENTS

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials.

CAUTION

Do not use in any finished feed (supplement, concentrate or complete feed) containing in excess of 2% bentonite.

FEEDING DIRECTIONS

Feed 40 to 100 g of tylosin per ton of complete feed for 2 to 6 weeks immediately after medicating with tylosin (250 mg tylosin per gallon) in drinking water for 3 to 10 days. Feed continuously as the sole ration.

MANUFACTURED BY:

BLUE BIRD FEED MILL
Any town, USA 12345

Net Weight lb (kg) on bag or bulk

^a Sourced from Tylan, NADA# 12-491

*The medicated feed label must state a single drug concentration.

