

# Tylosin Medicated Feed – Swine Ileitis

## Type C Medicated Feed

**CAUTION:** Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

### INDICATION

For control of porcine proliferative enteropathies (PPE, ileitis) associated with *Lawsonia intracellularis*

### ACTIVE DRUG INGREDIENT

Tylosin (as tylosin phosphate)<sup>a</sup> ..... 40 or 100 g/ton\*

### GUARANTEED ANALYSIS

Crude Protein, not less than.....	_____	%
Lysine, not less than.....	_____	%
Crude Fat, not less than .....	_____	%
Crude Fiber, not more than .....	_____	%
Calcium, not less than.....	_____	%
Calcium, not more than.....	_____	%
Phosphorus, not less than.....	_____	%
Salt <sup>1</sup> , not less than.....	_____	%
Salt <sup>1</sup> , not more than.....	_____	%
Sodium <sup>2</sup> , not less than .....	_____	%
Sodium <sup>2</sup> , not more than .....	_____	%
Selenium, not less than .....	_____	ppm
Zinc, not less than .....	_____	ppm

<sup>1</sup> If added.

<sup>2</sup> Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

### INGREDIENTS

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials.

### CAUTION

Do not use in any finished feed (supplement, concentrate or complete feed) containing in excess of 2% bentonite.

### FEEDING DIRECTIONS

Feed 100 g of tylosin per ton of complete feed for 21 days. Alternatively, feed 100 g of tylosin per ton of complete feed for at least three weeks, followed by 40 g tylosin per ton of complete feed until pigs reach market weight. Feed continuously as the sole ration.

### MANUFACTURED BY:

BLUE BIRD FEED MILL  
Any town, USA 12345

### Net Weight lb (kg) on bag or bulk

<sup>a</sup> Sourced from Tylan, NADA# 12-491

\*The medicated feed label must state a single drug concentration.

