

**Monensin Medicated Dairy Cattle Feed  
Type C Medicated Feed**

**For Total Mixed Rations**

**For Use in Dairy Cattle Feeds Only**

For Increased Milk Production Efficiency (production of marketable solids-corrected milk per unit of feed intake).

**Active Drug Ingredient**

Monensin, USP .....11 to 22 g/ton\*

**Guaranteed Analysis**

Crude Protein, not less than.....%  
Non-Protein Nitrogen (NPN)<sup>1</sup>, not more than.....%  
Crude Fat, not less than.....%  
Crude Fiber, not more than.....%  
Acid Detergent Fiber, not more than.....%  
Calcium, not less than.....%  
Calcium, not more than.....%  
Phosphorus, not less than.....%  
Salt<sup>2</sup>, not less than.....%  
Salt<sup>2</sup>, not more than.....%  
Sodium<sup>3</sup>, not less than.....%  
Sodium<sup>3</sup>, not more than.....%  
Potassium, not less than.....%  
Selenium, not less than.....ppm  
Vitamin A<sup>2,4</sup>, not less than.....I.U./lb

<sup>1</sup>When added.

<sup>2</sup> If added

<sup>3</sup>Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

<sup>4</sup>Other than precursors of Vitamin A.

**Ingredients**

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials.

**Feeding Directions**

Feed continuously to dry and lactating dairy cows a total mixed ration (“complete feed”) containing 11 to 22 g/ton monensin on a 100% dry matter basis.

\*Final printed label on formulated Type C medicated feed must bear a single drug concentration.

### **Caution**

Do not allow horses or other equines access to feed containing monensin. Ingestion of monensin by horses has been fatal. Monensin medicated cattle and goat feeds are safe for use in cattle and goats only. Consumption by unapproved species may result in toxic reactions. If feed refusals containing monensin are fed to other groups of cattle, the concentration of monensin in the refusals and amount of refusals fed should be taken into consideration to prevent monensin overdosing.

### **You May Notice**

- Reduced voluntary feed intake in dairy cows fed monensin. This reduction increases with higher doses of monensin fed. Rule out monensin as the cause of reduced feed intake before attributing to other causes such as illness, feed management, or the environment.
- Reduced milk fat percentage in dairy cows fed monensin. This reduction increases with higher doses of monensin fed.
- Increased incidence of cystic ovaries and metritis in dairy cows fed monensin.
- Reduced conception rates, increased services per animal, and extended days open and corresponding calving intervals in dairy cows fed monensin.

Have a comprehensive and ongoing nutritional, reproductive and herd health program in place when feeding monensin to dairy cows.

### **Warning**

A withdrawal time has not been established for pre-ruminating calves. Do not use in calves to be processed for veal.

**Manufactured By**  
Blue Bird Feed Mill  
Any town, USA 12345

Net Weight lb on bag or bulk