

Monensin Medicated Feedlot Cattle Feed

Type C Medicated Feed

**For Cattle Fed in Confinement for Slaughter
(For improved feed efficiency)**

For improved feed efficiency.

Active Drug Ingredient

Monensin, USP5 to 40 g/ton*
No additional improvement in feed efficiency has been shown from feeding
monensin at levels greater than 30 g/ton (360 mg monensin per head per day)

Guaranteed Analysis

Crude Protein, not less than %
Non-Protein Nitrogen (NPN)¹, not more than %
Crude Fat, not less than %
Crude Fiber, not more than %
Calcium, not less than %
Calcium, not more than %
Phosphorus, not less than %
Salt², not less than %
Salt², not more than %
Sodium³, not less than %
Sodium³, not more than %
Potassium, not less than %
Vitamin A^{2,4}, not less than I.U./lb

¹When added.

²If added.

³Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

⁴Other than precursors of Vitamin A.

Ingredients

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials.

* Final printed label on formulated Type C medicated feed must bear a single drug concentration.

Feeding Directions

Feed continuously in complete feed at a rate of 50 to 480 milligrams of monensin per head per day. No additional improvement in feed efficiency has been shown from feeding monensin at levels greater than 30 g/ton (360 mg monensin per head per day).

Caution

Do not allow horses or other equines access to feed containing monensin. Ingestion of monensin by horses has been fatal. Monensin medicated cattle and goat feeds are safe for use in cattle and goats only. Consumption by unapproved species may result in toxic reactions. Do not exceed the levels of monensin recommended in the feeding directions, as reduced average daily gains may result. If feed refusals containing monensin are fed to other groups of cattle, the concentration of monensin in the refusals and amount of refusals fed should be taken into consideration to prevent monensin overdosing.

Warning

A withdrawal period has not been established for this product in preruminating calves. Do not use in calves to be processed for veal.

Manufactured By
Blue Bird Feed Mill
Anytown, USA 12345

Net Weight lb on bag or bulk