

Net Weight lb (kg) on bag or bulk  
**Monensin Medicated Bobwhite Quail Feed**  
**Type C Medicated Feed**

For the prevention of coccidiosis caused by *Eimeria dispersa*, and *E. lettyae*.

**ACTIVE DRUG INGREDIENT**

Monensin, USP .....73 g/ton\*

**GUARANTEED ANALYSIS**

Crude Protein, not less than.....	_____	%
Lysine, not less than.....	_____	%
Methionine, not less than.....	_____	%
Crude Fat, not less than.....	_____	%
Crude Fiber, not more than.....	_____	%
Calcium, not less than.....	_____	%
Calcium, not more than.....	_____	%
Phosphorus, not less than.....	_____	%
Salt <sup>1</sup> , not less than.....	_____	%
Salt <sup>1</sup> , not more than.....	_____	%
Sodium <sup>2</sup> , not less than.....	_____	%
Sodium <sup>2</sup> , not more than.....	_____	%

<sup>1</sup>If added.

<sup>2</sup>Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

**INGREDIENTS**

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials.

**FEEDING DIRECTIONS**

Feed 73 g/ton of complete feed. Feed continuously as the sole ration.

**CAUTION**

Do not allow horses, other equines, mature turkeys, or guinea fowl access to feed containing monensin. Ingestion of monensin by horses and guinea fowl has been fatal.

**MANUFACTURED BY**

BLUE BIRD FEED MILL  
Any town, USA 12345

\*Final printed label on formulated Type C medicated feed must bear a single drug concentration.