

Net Weight lb (kg) on bag or bulk  
**Monensin Medicated Replacement Chicken Feed**  
**Type C Medicated Feed**

To aid in the prevention of coccidiosis caused by *Eimeria necatrix*, *E. tenella*, *E. acervulina*, *E. brunetti*, *E. mivati*, and *E. maxima*.

**ACTIVE DRUG INGREDIENT**

Monensin, USP .....90-110 g/ton\*

**GUARANTEED ANALYSIS**

Crude Protein, not less than.....%  
Lysine, not less than.....%  
Methionine, not less than.....%  
Crude Fat, not less than.....%  
Crude Fiber, not more than.....%  
Calcium, not less than.....%  
Calcium, not more than.....%  
Phosphorus, not less than.....%  
Salt<sup>1</sup>, not less than.....%  
Salt<sup>1</sup>, not more than.....%  
Sodium<sup>2</sup>, not less than.....%  
Sodium<sup>2</sup>, not more than.....%

<sup>1</sup>If added.

<sup>2</sup>Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

**INGREDIENTS**

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials.

**FEEDING DIRECTIONS**

Feed 90-110 g/ton of complete feed to replacement chickens intended for use as cage layers. Feed continuously as the sole ration.

**CAUTION**

Do not feed to laying chickens. Do not feed to chickens over 16 weeks of age. Do not allow horses, other equines, mature turkeys, or guinea fowl access to feed containing monensin. Ingestion of monensin by horses and guinea fowl has been fatal.

**MANUFACTURED BY**

BLUE BIRD FEED MILL  
Any town, USA 12345

\*Final printed label on formulated Type C medicated feed must bear a single drug concentration.