

Net weight lb (kg) on bag or bulk

Monensin/Tylosin phosphate Broiler Chicken Ration Type C Medicated Feed

As an aid in the prevention of coccidiosis caused by *Eimeria necatrix*, *E. tenella*, *E. acervulina*, *E. brunetti*, *E. mivati*, and *E. maxima*, and for increased rate of weight gain and improved feed efficiency in broiler chickens.

Active Drug Ingredient

Monensin.....90 to 110 g/ton
Tylosin phosphate.....4 to 50 g/ton

Guaranteed Analysis

Crude Protein, not less than..... %
Lysine, not less than..... %
Methionine, not less than..... %
Crude Fat, not less than..... %
Crude Fiber, not more than..... %
Calcium, not less than..... %
Calcium, not more than..... %
Phosphorus, not less than..... %
Salt¹, not less than..... %
Salt¹, not more than..... %
Sodium², not less than..... %
Sodium², not more than..... %

¹If added.

²Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

Ingredients

Each ingredient must be specifically named in accordance with the names and definitions adopted by the Association of American Feed Control Official's Publication.

Directions for Use

Feed continuously as sole ration.

(OVER)

CAUTION: Do not feed to laying chickens. Do not feed to chickens over 16 weeks of age. Do not allow horses, other equines, mature turkeys, or guinea fowl access to feed containing monensin. Ingestion of monensin by horses, mature turkeys, and guinea fowl has been fatal. Feed continuously as sole ration. In the absence of coccidiosis, the use of monensin with no withdrawal period may limit feed intake resulting in reduced weight gain.

MANUFACTURED BY

BLUE BIRD FEED MILL

Anytown, USA 12345