

Lot No. _____

BLUE BIRD TURKEY FEED

TYPE C MEDICATED FEED

For the prevention of coccidiosis caused by *E. adenoeides*, *E. meleagrimitis*, and *E. gallopavonis*, and as an aid in the control for transmissible enteritis in growing turkeys complicated by organisms susceptible to bacitracin methylene disalicylate.

ACTIVE INGREDIENT

Monensin (as monensin sodium)	54 to 90 g/ton
Bacitracin Methylene Disalicylate	200 g/ton

GUARANTEED ANALYSIS

Crude Protein, not less than	_____ %
Crude Fat, not less than	_____ %
Crude Fiber, not more than	_____ %

INGREDIENTS

Each ingredient must be specifically named (unless stated as such in the guaranteed analysis listing) in accordance with the names and definitions listed in Title 21 CFR 501.110.

DIRECTIONS FOR USE

Feeding Directions: Feed continuously as the sole ration. The optimum level depends upon the severity of coccidiosis.

CAUTION: Do not allow horses, other equines, mature turkeys or guinea fowl access to feed containing monensin. Ingestion of monensin by horses and guinea fowl has been fatal. Some strains of turkey coccidia may be monensin tolerant or resistant. Monensin may interfere with the development of immunity to turkey coccidiosis.

Store in a cool, dry place.

NET WEIGHT ON BAG OR BULK

Blue Bird Feed Mill
Robin, IN 12345