

**Monensin and Bacitracin Zinc Medicated Broiler Chicken Feed
(Feed Efficiency and Coccidiosis)
Type C Medicated Feed**

For improved feed efficiency; as an aid in the prevention of coccidiosis caused by *Eimeria necatrix*, *E. tenella*, *E. acervulina*, *E. brunetti*, *E. mivati*, and *E. maxima*.

ACTIVE DRUG INGREDIENTS

Monensin, USP..... 90 to 110 g/ton*
Bacitracin Zinc..... 10 to 30 g/ton*

GUARANTEED ANALYSIS

Crude Protein, not less than %
Lysine, not less than %
Methionine, not less than %
Crude Fat, not less than %
Crude Fiber, not more than %
Calcium, not less than %
Calcium, not more than %
Phosphorus, not less than %
Salt¹, not less than %
Salt¹, not more than %
Sodium², not less than %
Sodium², not more than %

¹If added

²Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

INGREDIENTS

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials.

FEEDING DIRECTIONS

Feed continuously as the sole ration.

CAUTION

Do not feed to laying chickens; in the absence of coccidiosis the use of monensin with no withdrawal period may limit feed intake resulting in reduced weight gain. Do not allow horses, other equines, mature turkeys or guinea fowl access to feed containing monensin. Ingestion of monensin by horses and guinea fowl has been fatal.

MANUFACTURED BY

BLUE BIRD FEED MILL, Any town, USA 12345

Net Weight lb (kg) on bag or bulk

*Final printed label on formulated Type C medicated feed must bear a single drug concentration.