

Net weight lb (kg) on bag or bulk

## Monensin/Bambermycins Growing Turkey Ration Type C Medicated Feed

For the prevention of coccidiosis caused by *Eimeria adenoeides*, *E. meleagrimitis*, and *E. gallopavonis*, and for increased rate of weight gain and improved feed efficiency in growing turkeys.

### ACTIVE DRUG INGREDIENT

Monensin.....	54 to 90 g/ton
Bambermycins.....	2 g/ton

### GUARANTEED ANALYSIS

Crude Protein, not less than.....	_____ %
Lysine, not less than.....	_____ %
Methionine, not less than.....	_____ %
Crude Fat, not less than.....	_____ %
Crude Fiber, not more than.....	_____ %
Calcium, not less than.....	_____ %
Calcium, not more than.....	_____ %
Phosphorus, not less than.....	_____ %
Salt <sup>1</sup> , not less than.....	_____ %
Salt <sup>1</sup> , not more than.....	_____ %
Sodium <sup>2</sup> , not less than.....	_____ %
Sodium <sup>2</sup> , not more than.....	_____ %

<sup>1</sup>If added.

<sup>2</sup>Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

### INGREDIENTS

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials.

### DIRECTIONS FOR USE

Feed continuously as the sole ration.

**CAUTION:** Do not allow horses, other equines, mature turkeys, or guinea fowl access to feed containing monensin. Ingestion of monensin by horses and guinea fowl has been fatal. Some strains of turkey coccidia may be monensin tolerant or resistant. Monensin may interfere with development of immunity to turkey coccidiosis.

**MANUFACTURED BY**  
BLUE BIRD FEED MILL  
Anytown, USA 12345

Net weight lb (kg) on bag or bulk  
**Monensin/Bambermycins Growing Turkey Ration**  
**Type C Medicated Feed**

For the prevention of coccidiosis caused by *Eimeria adenoeides*, *E. meleagrititis*, and *E. gallopavonis*, and for improved feed efficiency in growing turkeys.

**ACTIVE DRUG INGREDIENT**

Monensin.....54 to 90 g/ton  
Bambermycins.....1 to 2 g/ton

**GUARANTEED ANALYSIS**

Crude Protein, not less than.....                     %  
Lysine, not less than.....                     %  
Methionine, not less than.....                     %  
Crude Fat, not less than.....                     %  
Crude Fiber, not more than.....                     %  
Calcium, not less than.....                     %  
Calcium, not more than.....                     %  
Phosphorus, not less than.....                     %  
Salt<sup>1</sup>, not less than.....                     %  
Salt<sup>1</sup>, not more than.....                     %  
Sodium<sup>2</sup>, not less than.....                     %  
Sodium<sup>2</sup>, not more than.....                     %

<sup>1</sup>If added.

<sup>2</sup>Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

**INGREDIENTS**

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials.

**DIRECTIONS FOR USE**

Feed continuously as the sole ration.

**CAUTION:** Do not allow horses, other equines, mature turkeys, or guinea fowl access to feed containing monensin. Ingestion of monensin by horses and guinea fowl has been fatal. Some strains of turkey coccidia may be monensin tolerant or resistant. Monensin may interfere with development of immunity to turkey coccidiosis.

**MANUFACTURED BY**

BLUE BIRD FEED MILL  
Anytown, USA 12345