

Lot No. _____

**NET WEIGHT ON BAG OR BULK
BLUE BIRD TURKEY RATION
TYPE C MEDICATED FEED**

For the prevention of coccidiosis caused by *Eimeria adenoides*, *E. meleagrimitis*, and *E. gallopavonis*, and for increased rate of weight gain and improved feed efficiency in growing turkeys.

ACTIVE DRUG INGREDIENT

Monensin 54 to 90 g/ton
Virginiamycin 10 to 20 g/ton

GUARANTEED ANALYSIS

Crude Protein, not less than..... %
Lysine, not less than..... %
Methionine, not less than..... %
Crude Fat, not less than..... %
Crude Fiber, not more than..... %
Calcium, not less than..... %
Calcium, not more than..... %
Phosphorus, not less than..... %
Salt¹, not less than..... %
Salt¹, not more than..... %
Sodium², not less than..... %
Sodium², not more than..... %

¹If added.

²Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

INGREDIENTS

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials' Publication (AAFCO Official Publication).

DIRECTIONS FOR USE

Feed continuously as the sole ration from 1 day of age to 10 weeks of age.

CAUTION: Do not allow horses, other equines, mature turkeys or guinea fowl access to feed containing monensin. Ingestion of monensin by horses, mature turkeys, and guinea fowl has been fatal. Some strains of turkey coccidian may be monensin tolerant or resistant. Monensin may interfere with the development of immunity to turkey coccidiosis.

MANUFACTURED BY

Blue Bird Feed Mill
Anytown, USA 12345