

Net Weight lb (kg) on bag or bulk
Ractopamine Finishing Swine Feed Concentrate
Type B Medicated Feed
Do Not Feed Undiluted

IMPORTANT: MUST BE THOROUGHLY MIXED INTO FEED BEFORE USE

For increased rate of weight gain, improved feed efficiency and increased carcass leanness in finishing swine, weighing not less than 150 lbs, fed a complete ration containing at least 16% crude protein for the last 45 to 90 lbs of gain prior to slaughter.

ACTIVE DRUG INGREDIENT

Ractopamine hydrochloride..... 900 to 4920 g/ton*

GUARANTEED ANALYSIS

Crude Protein, not less than.....	_____ %
Lysine, not less than.....	_____ %
Crude Fat, not less than.....	_____ %
Crude Fiber, not more than.....	_____ %
Calcium, not less than.....	_____ %
Calcium, not more than.....	_____ %
Phosphorus, not less than.....	_____ %
Salt ¹ , not less than.....	_____ %
Salt ¹ , not more than.....	_____ %
Sodium ² , not less than.....	_____ %
Sodium ² , not more than.....	_____ %
Selenium, not less than.....	_____ ppm
Zinc, not less than.....	_____ ppm

¹If added.

²Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

INGREDIENTS

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials.

MIXING DIRECTIONS

Thoroughly mix ractopamine Type B Medicated Feed into one ton of complete swine feed, containing at least 16% crude protein, according to the table below to obtain the proper concentration in the Type C Medicated Feed. Prepare an intermediate pre-blend of the medicated feed prior to mixing in a complete feed. Thoroughly mix the required amount in a convenient quantity of feed ingredients, then add to the remaining feed ingredients to make a ton of complete feed. [Use only the portion of the table below on your Type B Medicated Feed product label that is applicable to the concentration of ractopamine in the Type B Medicated Feed which you manufacture.]

Concentration of Ractopamine In Type B Medicated Feed (grams/pound)	Pounds Type B Medicated Feed To Add Per Ton of Type C Medicated Feed	Resulting Ractopamine Concentration in Type C Medicated Feed ** (grams/ton)
0.45	10	4.5
	20	9
0.9	5	4.5
	10	9
1.35	3.3	4.5
	6.7	9
1.8	2.5	4.5
	5	9

** No increased benefit has been shown when ractopamine concentrations in the diet are greater than 4.5 g/ton (5 ppm)

CAUTION: Ractopamine may increase the number of injured and/or fatigued pigs during marketing. Not for use in breeding swine.

WARNING: The active ingredient in Paylean®, ractopamine hydrochloride, is a beta-adrenergic agonist. Individuals with cardiovascular disease should exercise special caution to avoid exposure. Not for use in humans. Keep out of the reach of children. The Paylean formulation (Type A Medicated Article) poses a low dust potential under usual conditions of handling and mixing. When mixing and handling Paylean, use protective clothing, impervious gloves, protective eye wear, and a NIOSH-approved dust mask. Operators should wash thoroughly with soap and water after handling. If accidental eye contact occurs, immediately rinse eyes thoroughly with water. If irritation persists, seek medical attention. The material safety data sheet contains more detailed occupational safety information. To report adverse effects, access medical information, or obtain additional product information, call 1-800-428-4441.

MANUFACTURED BY

BLUE BIRD FEED MILL

Any town, USA 12345

Paylean® is a registered trademark of Eli Lilly and Company.

*The final printed label must show only one drug level.