

# Narasin Medicated Swine Feed Type B Medicated Feed for Average Daily Gain

For increased rate of weight gain in growing-finishing swine when fed for at least 4 weeks.

## ACTIVE DRUG INGREDIENT

Narasin .....27.3 to 5400 g/ton\*

No increased benefit in rate of weight gain has been shown when narasin concentrations in the diet are greater than 13.6 g/ton.

## GUARANTEED ANALYSIS

Crude Protein, min.....	_____	%
Lysine, min.....	_____	%
Crude Fat, min .....	_____	%
Crude Fiber, max.....	_____	%
Calcium, min.....	_____	%
Calcium, max.....	_____	%
Phosphorus, min.....	_____	%
Salt <sup>a</sup> , min.....	_____	%
Salt <sup>a</sup> , max.....	_____	%
Sodium <sup>b</sup> , min.....	_____	%
Sodium <sup>b</sup> , max.....	_____	%
Selenium, min .....	_____	ppm
Zinc, min.....	_____	ppm

<sup>a</sup> Guarantee required only when nutrient added except when the feed is intended, represented or serves as a principal source of the nutrient.

<sup>b</sup> Sodium guarantee required only when total sodium exceeds that furnished by the maximum salt guarantee.

## INGREDIENTS

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials.

## MIXING DIRECTIONS

Thoroughly mix narasin Type B medicated feed with non-medicated swine feed according to the table below to obtain the proper concentration in the Type C Medicated Feed. Use only the portion of the table below on your Type B Medicated Feed product label that is applicable to the concentration of narasin in the Type B Medicated Feed which you manufacture. The following table gives examples of how some Type B Medicated Feed concentrations can be prepared.

\*Final printed label on formulated Type B medicated feed must bear a single drug concentration.

Narasin Concentration in Type B Feed (g/ton)	Desired Narasin Concentration in Type C Medicated Feed (g/ton)		
	13.6 g/ton	18.1 g/ton	27.2 g/ton
	lb of Type B per ton of Narasin Medicated Complete Feed		
100	272	362	544
500	54.4	72.4	108.8
1,000	27.2	36.2	54.4

### CAUTION

Do not allow adult turkeys, horses or other equines access to narasin formulations. Ingestion of narasin by these species has been fatal. Not approved for use in breeding animals because safety and effectiveness have not been evaluated in these animals.

Swine being fed with narasin should not have access to feeds containing pleuramutilins (e.g., tiamulin) as adverse reactions may occur. If signs of toxicity occur, discontinue use.

### WARNING



No withdrawal period is required when used according to the label.



Not for use in humans. When mixing and handling narasin, use protective clothing, impervious gloves and a dust mask. Operators should wash thoroughly with soap and water after handling. If accidental eye contact occurs, immediately rinse thoroughly with water. To report adverse effects, access medical information, or obtain additional product information, call 1-800-428-4441.

**Lot Number:** \_\_\_\_\_

**MANUFACTURED BY**  
**BLUE BIRD FEED MILL**  
 Any town, USA 12345

**Net Weight lb (kg) on bag or bulk**