

Narasin Medicated Swine Feed Type C Medicated Feed for Average Daily Gain

For increased rate of weight gain in growing-finishing swine when fed for at least 4 weeks.

ACTIVE DRUG INGREDIENT

Narasin 13.6 to 27.2 g/ton*

No increased benefit in rate of weight gain has been shown when narasin concentrations in the diet are greater than 13.6 g/ton.

GUARANTEED ANALYSIS

Crude Protein, min.....	_____	%
Lysine, min.....	_____	%
Crude Fat, min.....	_____	%
Crude Fiber, max.....	_____	%
Calcium, min.....	_____	%
Calcium, max.....	_____	%
Phosphorus, min.....	_____	%
Salt ^a , min.....	_____	%
Salt ^a , max.....	_____	%
Sodium ^b , min.....	_____	%
Sodium ^b , max.....	_____	%
Selenium, min.....	_____	ppm
Zinc, min.....	_____	ppm

^a Guarantee required only when nutrient added except when the feed is intended, represented or serves as a principal source of the nutrient.

^b Sodium guarantee required only when total sodium exceeds that furnished by the maximum salt guarantee.

INGREDIENTS

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials.

FEEDING DIRECTIONS

Feed continuously for at least four weeks to swine during the growing-finishing period as the sole ration.

*Final printed label on formulated Type C medicated feed must bear a single drug concentration.

CAUTION

Do not allow adult turkeys, horses or other equines access to narasin formulations. Ingestion of narasin by these species has been fatal. Not approved for use in breeding animals because safety and effectiveness have not been evaluated in these animals.

Swine being fed with narasin should not have access to feeds containing pleuramutilins (e.g., tiamulin) as adverse reactions may occur. If signs of toxicity occur, discontinue use.

Effectiveness has not been demonstrated when fed for durations less than four weeks.

WARNING



No withdrawal period is required when
used according to the label.



Not for use in humans. When mixing and handling narasin, use protective clothing, impervious gloves and a dust mask. Operators should wash thoroughly with soap and water after handling. If accidental eye contact occurs, immediately rinse thoroughly with water. To report adverse effects, access medical information, or obtain additional product information, call 1-800-428-4441.

Lot Number: _____

Manufactured On: _____

Type C Medicated Feed expires 2 months (8 weeks) from the "Manufacture On" date.

MANUFACTURED BY
BLUE BIRD FEED MILL
Any town, USA 12345

Net Weight lb (kg) on bag or bulk