

Net Weight lb (kg) on bag or bulk

# Tylosin Swine Dysentery Control Medicated Feed Type B Medicated Feed

**Do Not Feed Undiluted**

IMPORTANT: MUST BE THOROUGHLY MIXED INTO FEED BEFORE USE

For control of swine dysentery associated with *Brachyspira hyodysenteriae*.

## ACTIVE DRUG INGREDIENT

Tylosin phosphate ..... 101 to 20,000 g/ton\*

## GUARANTEED ANALYSIS

Crude Protein, not less than.....	_____%
Lysine, not less than.....	_____%
Crude Fat, not less than.....	_____%
Crude Fiber, not more than.....	_____%
Calcium, not less than.....	_____%
Calcium, not more than.....	_____%
Phosphorus, not less than.....	_____%
Salt <sup>1</sup> , not less than.....	_____%
Salt <sup>1</sup> , not more than.....	_____%
Sodium <sup>2</sup> , not less than.....	_____%
Sodium <sup>2</sup> , not more than.....	_____%
Selenium, not less than.....	_____ppm
Zinc, not less than.....	_____ppm

<sup>1</sup>If added.

<sup>2</sup>Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

## INGREDIENTS

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials.

## MIXING DIRECTIONS

Thoroughly mix tylosin Type B Medicated Feed into one ton of complete swine feed to obtain the correct concentration in the Type C Medicated Feed (Table 1).

**Table 1: Mixing Directions for Swine Medicated Feed**

Tylosin Concentration in Type B Feed (g/ton)	Desired Tylosin Concentration in Type C Medicated Feed (g/ton)	
	40	100
	lb of Type B per ton of Tylosin Medicated Complete Feed	
200	400	1000
2,000	40	100
20,000	4	10

**CAUTION**

Do not use in any finished feed (supplement, concentrate or complete feed) containing in excess of 2% bentonite.

**MANUFACTURED BY**

BLUE BIRD FEED MILL  
Any town, USA 12345

\*The medicated feed label must state a single drug concentration.