

Net Weight lb (kg) on bag or bulk

Tylosin Swine Dysentery Treatment and Control Medicated Feed Type C Medicated Feed

For treatment and control of swine dysentery associated with *Brachyspira hyodysenteriae* following initial medication with tylosin in drinking water.

ACTIVE DRUG INGREDIENT

Tylosin phosphate..... 40 to 100 g/ton*

GUARANTEED ANALYSIS

Crude Protein, not less than.....	_____	%
Lysine, not less than.....	_____	%
Crude Fat, not less than.....	_____	%
Crude Fiber, not more than.....	_____	%
Calcium, not less than.....	_____	%
Calcium, not more than.....	_____	%
Phosphorus, not less than.....	_____	%
Salt ¹ , not less than.....	_____	%
Salt ¹ , not more than.....	_____	%
Sodium ² , not less than.....	_____	%
Sodium ² , not more than.....	_____	%
Selenium, not less than.....	_____	ppm
Zinc, not less than.....	_____	ppm

¹If added.

²Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

INGREDIENTS

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials.

FEEDING DIRECTIONS

Administer in feed at 40 to 100 g of tylosin per ton of complete feed for two to six weeks after treatment with tylosin in drinking water (250 mg tylosin per gallon) for 3 to 10 days.

MANUFACTURED BY

BLUE BIRD FEED MILL
Any town, USA 12345

*The medicated feed label must state a single drug concentration.