

Net Weight lb (kg) on bag or bulk

Tylosin Swine Ileitis Control Medicated Feed Type C Medicated Feed

For control of porcine proliferative enteropathies (ileitis) associated with *Lawsonia intracellularis*.

ACTIVE DRUG INGREDIENT

Tylosin phosphate..... 40 or 100 g/ton*

GUARANTEED ANALYSIS

Crude Protein, not less than.....	_____	%
Lysine, not less than.....	_____	%
Crude Fat, not less than.....	_____	%
Crude Fiber, not more than.....	_____	%
Calcium, not less than.....	_____	%
Calcium, not more than.....	_____	%
Phosphorus, not less than.....	_____	%
Salt ¹ , not less than.....	_____	%
Salt ¹ , not more than.....	_____	%
Sodium ² , not less than.....	_____	%
Sodium ² , not more than.....	_____	%
Selenium, not less than.....	_____	ppm
Zinc, not less than.....	_____	ppm

¹If added.

²Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

INGREDIENTS

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials.

FEEDING DIRECTIONS

Feed 100 g of tylosin per ton of complete feed as the sole ration for 21 days. Alternatively, feed as the sole ration 100 g of tylosin per ton of complete feed for at least three weeks. Follow with 40 g tylosin per ton of complete feed until pigs reach market weight.

MANUFACTURED BY

BLUE BIRD FEED MILL
Any town, USA 12345

*The medicated feed label must state a single drug concentration.