

**Ractopamine and Tylosin Finishing Swine Medicated Feed
Type C Medicated Feed for
Treatment and Control of Swine Dysentery and Control of Ileitis**

For increased rate of weight gain, improved feed efficiency and increased carcass leanness in finishing swine, weighing not less than 150 lbs, fed a complete ration containing at least 16% crude protein for the last 45 to 90 lbs of gain prior to slaughter; for treatment and control of swine dysentery associated with *Brachyspira hyodysenteriae*; and for control of porcine proliferative enteropathies (PPE, ileitis) associated with *Lawsonia intracellularis*.

ACTIVE DRUG INGREDIENT

Ractopamine hydrochloride 4.5 to 9.0 g/ton*
Tylosin phosphate 40 to 100 g/ton**

GUARANTEED ANALYSIS

Crude Protein, not less than 16.00 %
Lysine, not less than _____ %
Crude Fat, not less than _____ %
Crude Fiber, not more than _____ %
Calcium, not less than _____ %
Calcium, not more than _____ %
Phosphorus, not less than _____ %
Salt¹, not less than _____ %
Salt¹, not more than _____ %
Sodium², not less than _____ %
Sodium², not more than _____ %
Selenium, not less than _____ ppm
Zinc, not less than _____ ppm

¹If added.

²Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

INGREDIENTS

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials.

FEEDING DIRECTIONS

Feed continuously as the sole ration to finishing swine weighing not less than 150 lbs for the last 45 to 90 lbs (group average) of weight gain prior to slaughter.

Feed 40 to 100 grams of tylosin per ton of complete feed for 2 to 6 weeks immediately after medicating with 250 mg tylosin tartrate (Tylan® Soluble) per gallon in drinking water for 3 to 10 days.

CAUTION

Ractopamine may increase the number of injured and/or fatigued pigs during marketing. Not for animals intended for breeding.

WARNING

The active ingredient in Paylean®, ractopamine hydrochloride, is a beta-adrenergic agonist. Individuals with cardiovascular disease should exercise special caution to avoid exposure. Not for use in humans. Keep out of the reach of children. The Paylean formulation (Type A Medicated Article) poses a low dust potential under usual conditions of handling and mixing. When mixing and handling Paylean, use protective clothing, impervious gloves, protective eye wear, and a NIOSH-approved dust mask. Operators should wash thoroughly with soap and water after handling. If accidental eye contact occurs, immediately rinse eyes thoroughly with water. If irritation persists, seek medical attention. The material safety data sheet contains more detailed occupational safety information. To report adverse effects, access medical information, or obtain additional product information, call 1-800-428-4441.

MANUFACTURED BY
BLUE BIRD FEED MILL
Any town, USA 12345

Net Weight lb (kg) on bag or bulk

Paylean® is a registered trademark of Eli Lilly and Company.

Tylan® is a registered trademark of Eli Lilly and Company.

*The medicated feed label must state a single drug concentration. No increased benefit has been shown when ractopamine concentrations in the diet are greater than 4.5 g/ton (5 ppm).

**The medicated feed label must state a single drug concentration.