



Tylan® (tylosin phosphate)

Sequential VFD ID Number

Veterinary Feed Directive (VFD) for use in Swine

Client: _____
Business _____
or Home _____
Address: _____
Phone #: _____

Veterinarian: _____
Address: _____
Phone #: _____

Approximate number of swine to be treated: _____ Location of animals: _____

Special Instructions and (or) Other Animal Identification (optional):

Indication, Drug Level in Medicated Feed, and Duration of Use (select one and specify the additional required information):

- For reduction in severity of effects of atrophic rhinitis. **Drug Level:** 100 grams per ton **Duration of Use:** _____ days
- For control of swine dysentery associated with *Brachyspira hyodysenteriae*.
Initial Drug Level: 100 grams per ton **Duration of Use:** _____ weeks (minimum of 3 weeks)
Subsequent Drug Level: 40 grams per ton **Duration of Use:** _____ weeks (until pigs reach market weight)
- For the treatment and control of swine dysentery associated with *Brachyspira hyodysenteriae* immediately after medicating with Tylan Soluble (tylosin) drinking water.
Drug Level: _____ grams per ton (40 to 100 g/ton) **Duration of Use:** _____ weeks (2 to 6 weeks)
- For control of porcine proliferative enteropathies (PPE, ileitis) associated with *Lawsonia intracellularis*.
Drug Level: 100 grams per ton **Duration of Use:** 21 days
- For control of porcine proliferative enteropathies (PPE, ileitis) associated with *Lawsonia intracellularis*.
Initial Drug Level: 100 grams per ton **Duration of Use:** _____ weeks (minimum of 3 weeks)
Subsequent Drug Level: 40 grams per ton **Duration of Use:** _____ weeks (until pigs reach market weight)
- For control of porcine proliferative enteropathies (PPE, ileitis) associated with *Lawsonia intracellularis* immediately after medicating with Tylan Soluble (tylosin) in drinking water.
Drug Level: _____ grams per ton (40 to 100 g/ton) **Duration of Use:** _____ weeks (2 to 6 weeks)

Use of feed containing this veterinary feed directive (VFD) drug in a manner other than as directed on the labeling (extra-label use) is not permitted.

Reorders (refills) are not permitted with this VFD drug.

WARNINGS

▶ **Residue Warning:** No withdrawal period is required. ◀

Combination Feeding with Other Drugs (select one):

- This VFD only authorizes the use of the VFD drug(s) cited in this order and is not intended to authorize the use of such drug(s) in combination with any other animal drugs.
- This VFD authorizes the use of the VFD drug(s) cited in this order in the following FDA-approved, conditionally approved, or indexed combination(s) in medicated feed that contains the VFD drug(s) as a component.

Drug(s) and Dose Range(s)		Specifications*
<input type="checkbox"/>	ractopamine hydrochloride at 4.5 to 9.0 g/ton** (supplied by Paylean™, NADA 141-172)	For use in finishing swine weighing not less than 150 lbs for the last 45 to 90 lbs (group average) of weight gain prior to slaughter.
<input type="checkbox"/>	Other FDA-approved, conditionally approved, or indexed combination:	

*for complete information see the approved Type C medicated feed label
**combination not available with atrophic rhinitis indication

- This VFD authorizes the use of the VFD drug(s) cited in this order in any FDA-approved, conditionally approved, or indexed combination(s) in medicated feed that contains the VFD drug(s) as a component.

VFD Issuance Date: _____

VFD Expiration Date: _____
(not to exceed 6 months from VFD issuance date)

Veterinarian's signature: _____

For technical service call: 1-800-428-4441
Elanco Animal Health a Division of Eli Lilly and Company
Indianapolis, IN 46285, USA

NADA 012-491, Approved by the FDA.

Tylan and the diagonal bar are trademarks owned or licensed by Eli Lilly and Company, its subsidiaries or affiliates.

White Copy (original) – Veterinarian

Canary Copy – Client

Pink Copy – Distribution