

Net Weight lb (kg) on bag or bulk

Tylosin Medicated Broiler Chicken Feed Type C Medicated Feed

To aid in the control of chronic respiratory disease caused by *Mycoplasma gallisepticum*.

ACTIVE DRUG INGREDIENT

Tylosin phosphate 800 to 1000 g/ton*

GUARANTEED ANALYSIS

Crude Protein, not less than.....	_____	%
Lysine, not less than.....	_____	%
Methionine, not less than.....	_____	%
Crude Fat, not less than.....	_____	%
Crude Fiber, not more than.....	_____	%
Calcium, not less than.....	_____	%
Calcium, not more than.....	_____	%
Phosphorus, not less than.....	_____	%
Salt ¹ , not less than.....	_____	%
Salt ¹ , not more than.....	_____	%
Sodium ² , not less than.....	_____	%
Sodium ² , not more than.....	_____	%

¹If added.

²Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

INGREDIENTS

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials.

FEEDING DIRECTIONS

Feed 800 to 1000 g/ton of complete feed. Administer in feed to chickens 0 to 5 days of age; follow with second administration in feed for 24 to 48 hours at 3 to 5 weeks of age.

WARNING

Withdraw 5 days before slaughter.

MANUFACTURED BY

BLUE BIRD FEED MILL
Any town, USA 12345

*The medicated feed label must state a single drug concentration.