

Net Weight lb (kg) on bag or bulk

**Tylosin Medicated Replacement Chicken Feed  
Type C Medicated Feed**

To aid in the control of chronic respiratory disease caused by *Mycoplasma gallisepticum*.

**ACTIVE DRUG INGREDIENT**

Tylosin phosphate ..... 1000 g/ton

**GUARANTEED ANALYSIS**

|  |       |   |
|--|-------|---|
| Crude Protein, not less than.....        | _____ | % |
| Lysine, not less than.....               | _____ | % |
| Methionine, not less than.....           | _____ | % |
| Crude Fat, not less than.....            | _____ | % |
| Crude Fiber, not more than.....          | _____ | % |
| Calcium, not less than.....              | _____ | % |
| Calcium, not more than.....              | _____ | % |
| Phosphorus, not less than.....           | _____ | % |
| Salt <sup>1</sup> , not less than.....   | _____ | % |
| Salt <sup>1</sup> , not more than.....   | _____ | % |
| Sodium <sup>2</sup> , not less than..... | _____ | % |
| Sodium <sup>2</sup> , not more than..... | _____ | % |

<sup>1</sup>If added.

<sup>2</sup>Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

**INGREDIENTS**

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials.

**FEEDING DIRECTIONS**

Feed 1000 g/ton of complete feed. Administer in feed to chickens 0 to 5 days of age; follow with second administration in feed for 24 to 48 hours at 3 to 5 weeks of age.

**WARNING**

Withdraw 5 days before slaughter.

**MANUFACTURED BY**

BLUE BIRD FEED MILL  
Any town, USA 12345