

Net Weight on Bag or Bulk
BLUE BIRD BROILER RATION
TYPE C MEDICATED FEED

For the prevention of coccidiosis caused by *Eimeria tenella*, *E. necatrix*, *E. acervulina*, *E. maxima*, *E. brunetti*, and *E. mivati*, and for increased rate of weight gain and improved feed efficiency in broiler chickens.

ACTIVE DRUG INGREDIENTS

Narasin.....	27 to 45 g/ton
Nicarbazin.....	27 to 45 g/ton
Bacitracin Methylene Disalicylate.....	4 to 50 g/ton

GUARANTEED ANALYSIS

Crude Protein, not less than.....	_____ %
Lysine, not less than.....	_____ %
Methionine, not less than.....	_____ %
Crude Fat, not less than.....	_____ %
Crude Fiber, not more than.....	_____ %
Calcium, not less than.....	_____ %
Calcium, not more than.....	_____ %
Phosphorus, not less than.....	_____ %
Salt ¹ , not less than.....	_____ %
Salt ¹ , not more than.....	_____ %
Sodium ² , not less than.....	_____ %
Sodium ² , not more than.....	_____ %

¹If added.

²Shall be guaranteed only when total sodium exceeds that furnished by the maximum salt guarantee.

INGREDIENTS

Each ingredient must be named in accordance with the names and definitions adopted by the Association of American Feed Control Officials' Publication (AAFCO Official Publication).

FEEDING DIRECTIONS

Feed continuously as the sole ration.

WARNING: Withdraw 5 days before slaughter.

CAUTION: For broiler chickens only. Do not feed to laying hens. Do not allow adult turkeys, horses or other equines access to formulations containing narasin. Ingestion of narasin by these species has been fatal. Nicarbazin medicated broilers may show reduced heat tolerance of exposed to high temperature and high humidity. Provide adequate drinking water and ventilation during these periods

MANUFACTURED BY
Blue Bird Feed Mills
Anytown, USA 12345

Lot Number _____