

**NET WEIGHT ON BAG OR BULK**

**BLUE BIRD BROILER RATION**

**TYPE C MEDICATED FEED**

For the prevention of coccidiosis caused by *Eimeria tenella*, *E. necatrix*, *E. acervulina*, *E. maxima*, *E. brunetti*, and *E. mivati*; and for increased rate of weight gain and improved feed efficiency in broiler chickens

**ACTIVE DRUG INGREDIENT**

Narasin.....27 – 45 g/ton  
Nicarbazin.....27 - 45 g/ton  
Lincomycin.....2 – 4 g/ton

**GUARANTEED ANALYSIS**

To provide nutritional requirements.

**INGREDIENTS**

As listed in the AAFCO Handbook.

**FEEDING DIRECTIONS**

Feed continuously as the only ration.

**WARNING:** Withdraw 5 days before slaughter.

**CAUTION:** Nicarbazin medicated broilers may show reduced heat tolerance if exposed to high temperature and high humidity. Provide adequate drinking water and ventilation during these periods. Do not allow adult turkeys, horses or other equines access to formulations containing narasin. Ingestion of narasin by these species has been fatal. Do not feed to laying hens. Do not allow rabbits, hamsters, guinea pigs, horses, or ruminants access to feeds containing lincomycin. Ingestion by these species may result in severe gastrointestinal effects.

**MANUFACTURED BY**

Blue Bird Feed Mills  
Anytown, USA 12345

Lot No. \_\_\_\_\_